

A Modern Introduction to Cardano and Ferrari Formulas in the Algebraic Equations

Kazuyuki FUJII *

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

We give a modern approach to the famous Cardano and Ferrari formulas in the algebraic equations with three and four degrees. Namely, we reconstruct these formulas from the point of view of superposition principle in quantum computation based on three and four level systems which are being developed by the author.

We also present a problem on some relation between Galois theory and Qudit theory.

*E-mail address : fujii@yokohama-cu.ac.jp

1 Introduction

The method to obtain algebraic solutions in (algebraic) equations has a very long history, see [1]. To solve a quadratic equation is very easy. To solve a cubic equation is not so easy and has been given by Cardano. How to solve a quartic equation is comparatively hard and Ferrari has given such solutions. After that many mathematicians including Lagrange, Gauss et al made how to solve equations refined. However, (algebraic) equations with degrees of more than four have in general no algebraic solutions, which was proved by two young mathematicians Abel and Galois.

By the way, we are studying a quantum computation based on multi-level systems of (laser-cooled) atoms. The three and four level systems are particularly interesting. In the theories the generalized Pauli and Walsh-Hadamard matrices (gates) play a central role. We don't mention the topics, see for example [2]–[8] as an introduction.

The aim of this note is to show that some fundamental techniques of the generalized Pauli and Walsh-Hadamard matrices are applicable to construct the Caldano and Ferrari formulas. The essential point is **the principle of superposition** which is the heart of Quantum Computation. Namely, we reconstruct the Caldano and Ferrari formulas from the view point of this principle.

Our method is so simple that it must have been known in some context. However, we cannot check all related references because the history is too long! We could not find such references within our effort. Anyway, by this note many students will be able to master the Caldano and Ferrari formulas without much effort.

2 Mathematical Preliminaries

We summarize the properties of Pauli matrices and Walsh-Hadamard matrix, and next state the corresponding ones of generalized Pauli matrices and Walsh-Hadamard matrix within our necessity.

Let $\{\sigma_1, \sigma_2, \sigma_3\}$ be Pauli matrices :

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1)$$

By (1) $\sigma_2 = i\sigma_1\sigma_3$, so that the essential elements of Pauli matrices are $\{\sigma_1, \sigma_3\}$ and they satisfy

$$\sigma_1^2 = \sigma_3^2 = \mathbf{1}_2 ; \quad \sigma_1^\dagger = \sigma_1, \quad \sigma_3^\dagger = \sigma_3 ; \quad \sigma_3\sigma_1 = -\sigma_1\sigma_3 = e^{i\pi}\sigma_1\sigma_3. \quad (2)$$

The Walsh–Hadamard matrix is defined by

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \in O(2) \subset U(2). \quad (3)$$

This matrix (or transformation) is unitary and it plays a very important role in Quantum Computation. Moreover it is easy to realize it in Quantum Optics as shown in [5]. Let us list some important properties of W :

$$W^2 = \mathbf{1}_2, \quad W^\dagger = W = W^{-1}, \quad (4)$$

$$\sigma_1 = W\sigma_3W^{-1}, \quad (5)$$

The check is very easy.

Next let us generalize Pauli matrices to higher dimensional cases. Let $\{\Sigma_1, \Sigma_3\}$ be the following matrices in $M(n, \mathbf{C})$

$$\Sigma_1 = \begin{pmatrix} 0 & & & & 1 \\ 1 & 0 & & & \\ & 1 & 0 & & \\ & & 1 & \cdot & \\ & & & \cdot & \cdot \\ & & & & 1 & 0 \end{pmatrix}, \quad \Sigma_3 = \begin{pmatrix} 1 & & & & \\ & \sigma & & & \\ & & \sigma^2 & & \\ & & & \cdot & \\ & & & & \cdot \\ & & & & & \sigma^{n-1} \end{pmatrix} \quad (6)$$

where σ is a primitive root of unity $\sigma^n = 1$ ($\sigma = e^{\frac{2\pi i}{n}}$). We note that

$$\bar{\sigma} = \sigma^{n-1}, \quad 1 + \sigma + \cdots + \sigma^{n-1} = 0.$$

The two matrices $\{\Sigma_1, \Sigma_3\}$ are generalizations of Pauli matrices $\{\sigma_1, \sigma_3\}$, but they are not hermitian. Here we list some of their important properties :

$$\Sigma_1^n = \Sigma_3^n = \mathbf{1}_n ; \quad \Sigma_1^\dagger = \Sigma_1^{n-1}, \quad \Sigma_3^\dagger = \Sigma_3^{n-1} ; \quad \Sigma_3 \Sigma_1 = \sigma \Sigma_1 \Sigma_3 . \quad (7)$$

For $n = 3$ and $n = 4$ Σ_1 and its powers are given respectively as

$$\Sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \\ & 1 & 0 \end{pmatrix}, \quad \Sigma_1^2 = \begin{pmatrix} 0 & 1 & \\ & 0 & 1 \\ 1 & & 0 \end{pmatrix} \quad (8)$$

and

$$\Sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \\ & 1 & 0 \\ & & 1 & 0 \end{pmatrix}, \quad \Sigma_1^2 = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ 1 & & 0 & \\ & 1 & & 0 \end{pmatrix}, \quad \Sigma_1^3 = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ 1 & & & 0 \end{pmatrix}. \quad (9)$$

If we define a Vandermonde matrix W based on σ as

$$W = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdot & \cdot & \cdot & 1 \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdot & \cdot & \cdot & \sigma^{(n-1)^2} \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdot & \cdot & \cdot & \sigma^{(n-1)(n-2)} \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ 1 & \sigma^2 & \sigma^4 & \cdot & \cdot & \cdot & \sigma^{2(n-1)} \\ 1 & \sigma & \sigma^2 & \cdot & \cdot & \cdot & \sigma^{n-1} \end{pmatrix}, \quad (10)$$

$$W^\dagger = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdot & \cdot & \cdot & 1 \\ 1 & \sigma & \sigma^2 & \cdot & \cdot & \cdot & \sigma^{n-1} \\ 1 & \sigma^2 & \sigma^4 & \cdot & \cdot & \cdot & \sigma^{2(n-1)} \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdot & \cdot & \cdot & \sigma^{(n-1)(n-2)} \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdot & \cdot & \cdot & \sigma^{(n-1)^2} \end{pmatrix}, \quad (11)$$

then it is not difficult to see

$$W^\dagger W = WW^\dagger = \mathbf{1}_n, \quad \Sigma_1 = W\Sigma_3W^\dagger = W\Sigma_3W^{-1}. \quad (12)$$

Since W corresponds to the Walsh–Hadamard matrix (3), so it may be possible to call W the generalized Walsh–Hadamard matrix. If we write $W^\dagger = (w_{ab})$, then

$$w_{ab} = \frac{1}{\sqrt{n}}\sigma^{ab} = \frac{1}{\sqrt{n}}\exp\left(\frac{2\pi i}{n}ab\right) \quad \text{for } 0 \leq a, b \leq n-1.$$

This is just the coefficient matrix of Discrete Fourier Transform (DFT) if $n = 2^k$ for some $k \in \mathbf{N}$, see [9].

For $n = 3$ and $n = 4$ W is given respectively as

$$W = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \frac{-1-i\sqrt{3}}{2} & \frac{-1+i\sqrt{3}}{2} \\ 1 & \frac{-1+i\sqrt{3}}{2} & \frac{-1-i\sqrt{3}}{2} \end{pmatrix} \quad (13)$$

and

$$W = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \sigma^3 & \sigma^2 & \sigma \\ 1 & \sigma^2 & 1 & \sigma^2 \\ 1 & \sigma & \sigma^2 & \sigma^3 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -i & -1 & i \\ 1 & -1 & 1 & -1 \\ 1 & i & -1 & -i \end{pmatrix}. \quad (14)$$

We note that the generalized Pauli and Walsh–Hadamard matrices in three and four level systems can be constructed (by using Rabi oscillations of several types) in a quantum optical manner, see [5] and [6].

3 Cardano Formula

We consider an algebraic equation with three degrees

$$x^3 + ax^2 + bx + c = 0. \quad (15)$$

Then by replacing $x \longrightarrow x - (a/3)$ we have

$$x^3 + px + q = 0$$

with $p = b - \frac{a^2}{3}$ and $q = c - \frac{ab}{3} + \frac{2a^3}{27}$. However, we usually set

$$x^3 + 3px + q = 0 \quad (16)$$

with

$$p = \frac{b}{3} - \frac{a^2}{9}, \quad q = c - \frac{ab}{3} + \frac{2a^3}{27}.$$

For this equation we have the famous solutions called **Cardano formula** : Three solutions are given by

$$x = u_0 + v_0, \quad \sigma u_0 + \sigma^2 v_0, \quad \sigma^2 u_0 + \sigma v_0, \quad (17)$$

where u_0 and v_0 are primitive solutions of binomial equations

$$u^3 = \frac{-q + \sqrt{q^2 + 4p^3}}{2}, \quad v^3 = \frac{-q - \sqrt{q^2 + 4p^3}}{2}$$

and $\sigma = e^{\frac{2\pi i}{3}}$.

The solutions (17) look like (classical) superpositions of u_0 and v_0 . From here we reconstruct the solutions by using generalized Pauli and Walsh–Hadamard matrices (8), (13) in three level systems, which makes the principle of superposition more clearly.

Let x_1, x_2, x_3 be three solutions of (16) and set

$$A = \begin{pmatrix} x_1 & & \\ & x_2 & \\ & & x_3 \end{pmatrix}, \quad x_j^3 + 3px_j + q = 0. \quad (18)$$

Remind that $x_1 + x_2 + x_3 = 0$ from (16), so $\text{tr} A = 0$. Then we have

$$A^3 + 3pA + qE = 0, \quad (19)$$

where E is a unit matrix. Now we consider superpositions of solutions x_j ($j = 1, 2, 3$).

For that we operate W in (13) on (19) as the adjoint action ($W^\dagger W = E$)

$$(W^\dagger A W)^3 + 3p(W^\dagger A W) + qE = 0. \quad (20)$$

Let us calculate $W^\dagger AW$:

$$\begin{aligned}
W^\dagger AW &= \frac{1}{3} \begin{pmatrix} x_1 + x_2 + x_3 & x_1 + \sigma^2 x_2 + \sigma x_3 & x_1 + \sigma x_2 + \sigma^2 x_3 \\ x_1 + \sigma x_2 + \sigma^2 x_3 & x_1 + x_2 + x_3 & x_1 + \sigma^2 x_2 + \sigma x_3 \\ x_1 + \sigma^2 x_2 + \sigma x_3 & x_1 + \sigma x_2 + \sigma^2 x_3 & x_1 + x_2 + x_3 \end{pmatrix} \\
&\equiv \begin{pmatrix} 0 & \beta & \alpha \\ \alpha & 0 & \beta \\ \beta & \alpha & 0 \end{pmatrix} = \alpha \begin{pmatrix} 0 & 1 \\ 1 & 0 \\ 1 & 0 \end{pmatrix} + \beta \begin{pmatrix} 0 & 1 \\ 0 & 1 \\ 1 & 0 \end{pmatrix} \\
&= \alpha \Sigma_1 + \beta \Sigma_1^2.
\end{aligned} \tag{21}$$

By the way,

$$(0, \beta, \alpha) = (x_1, x_2, x_3) \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix}.$$

Conversely,

$$(x_1, x_2, x_3) = (0, \beta, \alpha) \left\{ \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} \right\}^{-1} = (0, \beta, \alpha) \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix}.$$

Therefore we have

$$x_1 = \alpha + \beta, \quad x_2 = \sigma^2 \alpha + \sigma \beta, \quad x_3 = \sigma \alpha + \sigma^2 \beta. \tag{22}$$

Compare this with (17). Let us solve (20).

$$\begin{aligned}
0 &= (\alpha \Sigma_1 + \beta \Sigma_1^2)^3 + 3p(\alpha \Sigma_1 + \beta \Sigma_1^2) + qE \\
&= (\alpha^3 + \beta^3 + q)E + 3\alpha(\alpha\beta + p)\Sigma_1 + 3\beta(\alpha\beta + p)\Sigma_1^2
\end{aligned} \tag{23}$$

Therefore

$$\begin{cases} \alpha^3 + \beta^3 = -q \\ \alpha\beta = -p \end{cases} \tag{24}$$

Since $\beta = -p/\alpha$, we have

$$\alpha^3 - \frac{p^3}{\alpha^3} = -q \implies \alpha^6 + q\alpha^3 - p^3 = 0.$$

If we set $t = \alpha^3$, then $t^2 + qt - p^3 = 0$, the solutions are just

$$t = \frac{-q + \sqrt{q^2 + 4p^3}}{2}, \quad \frac{-q - \sqrt{q^2 + 4p^3}}{2}. \quad (25)$$

Since α and β are symmetric in the equation (24), α and β are primitive solutions of equations

$$\alpha^3 = \frac{-q + \sqrt{q^2 + 4p^3}}{2}, \quad \beta^3 = \frac{-q - \sqrt{q^2 + 4p^3}}{2}. \quad (26)$$

This is our derivation of **Cardano formula**.

4 Ferrari Formula

In this section we consider an algebraic equation with four degrees. By the same reasoning in the preceding section we have only to treat

$$x^4 + px^2 + qx + r = 0 \quad (27)$$

For this equation we have the famous solutions called **Ferrari formula** : Let λ_0 be a solution of the (decomposition) equation with three degrees

$$\lambda^3 - p\lambda^2 - 4r\lambda + (4pr - q^2) = 0. \quad (28)$$

Then four solutions of (27) are given by ones of following two equations with two degrees

$$x^2 \pm \sqrt{\lambda_0 - p} \left\{ x - \frac{q}{2(\lambda_0 - p)} \right\} + \frac{\lambda_0}{2} = 0. \quad (29)$$

This formula is not similar to that of Cardano. From this formula we cannot see the principle of superposition (of solutions), so we are dissatisfied at this one. From here we reconstruct the solutions by using generalized Pauli and Walsh–Hadamard matrices (9), (14) in four level systems to make the principle of superposition clear. Our method may be fresh to not only students (non–experts) but also mathematicians.

Let x_1, x_2, x_3, x_4 be four solutions of (27) and set

$$A = \begin{pmatrix} x_1 & & & \\ & x_2 & & \\ & & x_3 & \\ & & & x_4 \end{pmatrix}, \quad x_j^4 + px_j^2 + qx_j + r = 0. \quad (30)$$

Remind that $x_1 + x_2 + x_3 + x_4 = 0$ from (27), so $\text{tr}A = 0$. Then we have

$$A^4 + pA^2 + qA + rE = 0, \quad (31)$$

where E is a unit matrix. Similarly in the preceding section we consider superpositions of solutions x_j ($j = 1, 2, 3, 4$). We operate W in (14) on (31) as the adjoint action ($W^\dagger W = E$)

$$(W^\dagger AW)^4 + p(W^\dagger AW)^2 + qW^\dagger AW + rE = 0. \quad (32)$$

The calculation of $W^\dagger AW$ is as follows :

$$\begin{aligned} & W^\dagger AW \\ &= \begin{pmatrix} 0 & \beta & \gamma & \alpha \\ \alpha & 0 & \beta & \gamma \\ \gamma & \alpha & 0 & \beta \\ \beta & \gamma & \alpha & 0 \end{pmatrix} = \alpha \begin{pmatrix} 0 & & & 1 \\ 1 & 0 & & \\ & 1 & 0 & \\ & & 1 & 0 \end{pmatrix} + \gamma \begin{pmatrix} 0 & & 1 & \\ & 0 & & 1 \\ 1 & & 0 & \\ & 1 & & 0 \end{pmatrix} + \beta \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & 0 & 1 \\ 1 & & & 0 \end{pmatrix} \\ &= \alpha \Sigma_1 + \gamma \Sigma_1^2 + \beta \Sigma_1^3, \end{aligned} \quad (33)$$

where

$$\begin{aligned} 0 &= \frac{1}{4}(x_1 + x_2 + x_3 + x_4), \quad \alpha = \frac{1}{4}(x_1 + \sigma x_2 + \sigma^2 x_3 + \sigma^3 x_4), \\ \gamma &= \frac{1}{4}(x_1 + \sigma^2 x_2 + x_3 + \sigma^2 x_4), \quad \beta = \frac{1}{4}(x_1 + \sigma^3 x_2 + \sigma^2 x_3 + \sigma x_4) \end{aligned} \quad (34)$$

and $\sigma = e^{\frac{2\pi i}{4}} = i$. Then

$$(0, \beta, \gamma, \alpha) = (x_1, x_2, x_3, x_4) \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \sigma^3 & \sigma^2 & \sigma \\ 1 & \sigma^2 & 1 & \sigma^2 \\ 1 & \sigma & \sigma^2 & \sigma^3 \end{pmatrix}.$$

Conversely,

$$\begin{aligned} & (x_1, x_2, x_3, x_4) \\ & = (0, \beta, \gamma, \alpha) \left\{ \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \sigma^3 & \sigma^2 & \sigma \\ 1 & \sigma^2 & 1 & \sigma^2 \\ 1 & \sigma & \sigma^2 & \sigma^3 \end{pmatrix} \right\}^{-1} = (0, \beta, \gamma, \alpha) \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 & \sigma^3 \\ 1 & \sigma^2 & 1 & \sigma^2 \\ 1 & \sigma^3 & \sigma^2 & \sigma \end{pmatrix}, \end{aligned}$$

so we have

$$\begin{aligned} x_1 &= \alpha + \gamma + \beta, & x_2 &= \sigma^3 \alpha + \sigma^2 \gamma + \sigma \beta, \\ x_3 &= \sigma^2 \alpha + \gamma + \sigma^2 \beta, & x_4 &= \sigma \alpha + \sigma^2 \gamma + \sigma^3 \beta. \end{aligned} \quad (35)$$

This is just superpositions of α , γ and β . Next let us determine them. From (33)

$$(W^\dagger AW)^2 = (\gamma^2 + 2\alpha\beta)E + 2\gamma\beta\Sigma_1 + (\alpha^2 + \beta^2)\Sigma_1^2 + 2\alpha\gamma\Sigma_1^3$$

and

$$\begin{aligned} (W^\dagger AW)^4 &= \{(W^\dagger AW)^2\}^2 = \{(\alpha^2 + \beta^2)^2 + (\gamma^2 + 2\alpha\beta)^2 + 8\alpha\beta\gamma^2\} E + \\ & 4\{(\alpha^2 + \beta^2)\alpha + (\gamma^2 + 2\alpha\beta)\beta\} \gamma \Sigma_1 + \\ & 2(\alpha^2 + \beta^2)(3\gamma^2 + 2\alpha\beta)\Sigma_1^2 + \\ & 4\{(\alpha^2 + \beta^2)\beta + (\gamma^2 + 2\alpha\beta)\alpha\} \gamma \Sigma_1^3. \end{aligned}$$

Then the equation (32) gives

$$\begin{cases} (\alpha^2 + \beta^2)^2 + (\gamma^2 + 2\alpha\beta)^2 + 8\alpha\beta\gamma^2 + p(\gamma^2 + 2\alpha\beta) + r = 0 \\ 4\{(\alpha^2 + \beta^2)\alpha + (\gamma^2 + 2\alpha\beta)\beta\} \gamma + 2p\gamma\beta + q\alpha = 0 \\ 2(\alpha^2 + \beta^2)(3\gamma^2 + 2\alpha\beta) + p(\alpha^2 + \beta^2) + q\gamma = 0 \\ 4\{(\alpha^2 + \beta^2)\beta + (\gamma^2 + 2\alpha\beta)\alpha\} \gamma + 2p\alpha\gamma + q\beta = 0 \end{cases} \quad (36)$$

Now let us transform the equations into a more convenient form (this point is important).

$$\begin{cases} (a) & (\alpha^2 + \beta^2)^2 + (\gamma^2 + 2\alpha\beta)^2 + 8\alpha\beta\gamma^2 + p(\gamma^2 + 2\alpha\beta) + r = 0 \\ (b) & 2\gamma\beta(2\gamma^2 + 6\alpha\beta + p) + 4\alpha^3\gamma + q\alpha = 0 \\ (c) & (\alpha^2 + \beta^2)(6\gamma^2 + 4\alpha\beta + p) + q\gamma = 0 \\ (d) & 2\alpha\gamma(2\gamma^2 + 6\alpha\beta + p) + 4\beta^3\gamma + q\beta = 0 \end{cases} \quad (37)$$

From (b) and (d)

$$-\frac{4\alpha^3\gamma + q\alpha}{2\gamma\beta} = 2\gamma^2 + 6\alpha\beta + p = -\frac{4\beta^3\gamma + q\beta}{2\alpha\gamma},$$

so

$$4\alpha^4\gamma + q\alpha^2 = 4\beta^4\gamma + q\beta^2 \implies (\alpha^2 - \beta^2) \{4(\alpha^2 + \beta^2)\gamma + q\} = 0,$$

so we have

$$\alpha^2 + \beta^2 = -\frac{q}{4\gamma}. \quad (38)$$

Substituting this into (c)

$$6\gamma^2 + 4\alpha\beta + p - 4\gamma^2 = 0 \implies \gamma^2 + 2\alpha\beta = -\frac{p}{2} \quad \text{or} \quad 2\alpha\beta = -\gamma^2 - \frac{p}{2}. \quad (39)$$

Substituting these into (a)

$$\left(-\frac{q}{4\gamma}\right)^2 + \left(-\frac{p}{2}\right)^2 - 4\left(\gamma^2 + \frac{p}{2}\right)\gamma^2 - \frac{p^2}{2} + r = 0.$$

Rearranging this equation

$$64\gamma^6 + 32p\gamma^4 - 4(4r - p^2)\gamma^2 - q^2 = 0.$$

If we set $\Gamma = \gamma^2$, then we obtain the algebraic equation with three degrees

$$64\Gamma^3 + 32p\Gamma^2 - 4(4r - p^2)\Gamma - q^2 = 0. \quad (40)$$

Since this equation can be solved (in the preceding section), we obtain $\gamma = \sqrt{\Gamma}$ for the largest real solution Γ . Next we look for α and β . From (38) and (39)

$$\begin{cases} \alpha^2 + \beta^2 = -\frac{q}{4\gamma} \\ \alpha\beta = -\frac{1}{2}\left(\Gamma + \frac{p}{2}\right) \end{cases} \quad (41)$$

By the same method in (24) α and β are primitive solutions of equations

$$\alpha^2 = \frac{1}{2} \left\{ -\frac{q}{4\gamma} + \sqrt{\left(\frac{q}{4\gamma}\right)^2 - \left(\Gamma + \frac{p}{2}\right)^2} \right\}, \quad (42)$$

$$\beta^2 = \frac{1}{2} \left\{ -\frac{q}{4\gamma} - \sqrt{\left(\frac{q}{4\gamma}\right)^2 - \left(\Gamma + \frac{p}{2}\right)^2} \right\}. \quad (43)$$

Namely, we determined α , β and γ completely.

This is our derivation (a variation) of **Ferrari formula**.

5 A (Real) Problem

In this section let us consider an algebraic equation with five degrees.

$$x^5 + px^3 + qx^2 + rx + s = 0. \quad (44)$$

We can apply the same method in the case of three and four degrees to this equation, however we cannot solve this one in an algebraic manner as in the preceding sections (Abel and Galois). We know the Galois theory to decide whether an algebraic equation has algebraic solutions or not. This is a classical theory.

Then we have some natural questions : Is there some relation between the Galois theory and the principle of superposition ? What is a role of Galois theory in general Qudit theory ? Moreover, is it possible to consider a quantum Galois theory in general Qudit theory ?

These are interesting problems in Quantum Computation.

Acknowledgment.

The author wishes to thank Shin'ichi Nojiri for his helpful comments and suggestions.

Appendix Euler Formula

Akira Asada pointed out to me that my method to derive the variation of Ferrari formula was similar to that of Euler which I didn't know when writing this note. In fact his point was correct, so let us copy the Euler formula in [10]¹ for the completeness (this formula is not necessarily popular even in Mathematics).

For the quartic equation (27) we consider the cubic equation

$$t^3 + \frac{p}{2}t^2 + \left(\frac{p^2}{16} - \frac{r}{4}\right)t - \frac{q^2}{64} = 0. \quad (45)$$

¹I could not find a textbook written in English which comments on the formula

We note that this equation is just our (reduction) equation (40). Since this equation has three solutions (t_1, t_2, t_3) , we choose signs of square roots $\sqrt{t_1}, \sqrt{t_2}, \sqrt{t_3}$ to satisfy

$$\sqrt{t_1}\sqrt{t_2}\sqrt{t_3} = -\frac{q}{8}. \quad (46)$$

Then the solutions are given by

$$\begin{aligned} x_1 &= \sqrt{t_1} + \sqrt{t_2} + \sqrt{t_3}, & x_2 &= \sqrt{t_1} - \sqrt{t_2} - \sqrt{t_3}, \\ x_3 &= -\sqrt{t_1} + \sqrt{t_2} - \sqrt{t_3}, & x_4 &= -\sqrt{t_1} - \sqrt{t_2} + \sqrt{t_3}. \end{aligned} \quad (47)$$

To derive the solutions one use various relations between solutions (roots) and coefficients, which is quite complicated.

Our method is just a modern derivation of the Cardano and Euler formulas which doesn't use relations between solutions (roots) and coefficients in an apparent manner.

References

- [1] Mathematical Society of Japan : Encyclopedic Dictionary of Mathematics, MIT Press, 1987.
- [2] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, J. Applied Math, 2(2002), 371, quant-ph/0103011.
- [3] K. Fujii : Exchange Gate on the Qudit Space and Fock Space, J. Opt. B : Quantum Semiclass. Opt, 5(2003), S613, quant-ph/0207002.
- [4] K. Fujii : How To Treat An N-Level System : A Proposal, quant-ph/0302050.
- [5] K. Fujii : Quantum Optical Construction of Generalized Pauli and Walsh-Hadamard Matrices in Three Level Systems, quant-ph/0309132.
- [6] K. Fujii : Quantum Optical Construction of Generalized Walsh-Hadamard Matrix in Four Level Systems, in preparation.

- [7] K. Fujii, K. Higashida, R. Kato and Y. Wada : N Level System with RWA and Analytical Solutions Revisited, quant-ph/0307066.
- [8] K. Funahashi : Explicit Construction of Controlled-U and Unitary Transformation in Two-Qudit, quant-ph/0304078.
- [9] P. W. Shor : Polynomial-Time Algorithms for Prime Factorization and Discrete Logarithms on a Quantum Computer, SIAM J. Sci. Statist. Comput. 26(1997) 1484, quant-ph/9508027.
- [10] I. Yakabe : Introduction to Galois Theory (in Japanese), Gendaisugakusha, Kyoto, 1976.